BLOOD COLLECTION TUBE GUIDE - ORDER OF DRAW

This Document is Uncontrolled in Printed Form



TUBE GUIDE ORDER OF DRAW

Special Instructions

- ALL tubes must be labelled with patient's full name, date of birth, date and time of collection, and collector's signature.
- Never inject blood into the tube through a needle
 Fill tubes to indicator line and mix all samples.
- Fill tubes to indicator line and mix all samples immediately by gently inverting 8 10 times

Tube Content and Volume	Test list					
	Adult collection – 2 each of: Aerobic (green cap) & Anaerobic (orange cap), each filled with 8-10 mls blood Paediatric collection – 1 only of BacT/ALERT PF Plus (yellow cap) – up to 4 mls.					
BLOOD CULTURE BOTTLES Adult = Green and Orange Paed = Yellow	Store and transport at room temperature.					
	Cell markers/lymphocyte surface markers/flow cytometry/CD4 and CD8 HLA Tissue Typing					
PALE YELLOW ACD 9ml	Collect Monday - Thursday ONLY. Samples must reach the processing department before 2pm					
	All coagulation tests: APTT, Fibrinogen, INR, Coagulation Factor assays, VWF, FVIII and DDimer and Coagulation Profile (Note Coagulation profile also requires 4 ml EDTA)					
BLUE Sodium Citrate 2.7ml	□ D-Dimers - URGENT transport to laboratory required					
Sodium Citrate 2.7ml	Correct fill volume is Critical for analysis – tubes must be filled to the line indicated on the tube					
	General Biochemistry, hormones, troponin, lipids, iron, immunology, serology and therapeutic drug levels					
GOLD	☐ Troponin T - URGENT transport to laboratory required ☐					
Gel-SST 8.5ml	For ALL Drug levels – note time of last dose					
GREEN Lithium Heparin 6ml	Chromosomes Studies, Karyotyping Collect Monday - Thursday ONLY. Samples must reach the processing department before 2 pm					
PURPLE/LAVENDER EDTA 4ml	FBE/FBC, Diff, ESR, Hb, Platelets, Reticulocytes, IM Screen – all in 1 x 4ml sample Note an additional dedicated 2-4ml tube is required for each of: Cyclosporin, Everolimus, Sirolimus, Tacrolimus (note time of last dose) Haemochromatosis, Fragile X, FVL/PGM, Malaria, HbA1c, HLA B 27, HLA DQ 2/8, Kleihauer screen(FMH), Hb electrophoresis.					
	⇔ Ammonia – URGENT testing required, must reach lab within 20 minutes Blood Bank Testing: Antenatal Blood Group & Antibody Screen, Blood Group, Cross Match, Direct Coombs, Group					
	& Hold, Cold Agglutinins, Red Cell Antibody Screen All requests forms and tubes must signed by the collector, the labelled tubes must match the request forms, and include patients full name,					
PINK EDTA 6ml	DOB, hospital number, date and time of collection.					
	Aldosterone, renin, ACTH					
EDTA PEARL EDTA with gel	Samples must be collected before 10 am. These tests are restricted to collection centres with dedicated equipment – samples must be centrifuged before transport.					
NAVY BLUE Trace Metal Free 6ml	Trace Metal Analysis: Arsenic, Cadmium, Chromium, Cobalt, Lead, Mercury, Red Cell Magnesium, Zinc					
•	Glucose, Blood Alcohol, Lactate					
GREY Fluoride Oxalate 4ml	□ Lactate – URGENT testing required, must reach lab within 60 minutes					
	Immunoserology for TB – Quantiferon testing					
QUANTIFERON GOLD KIT 1 ml each	Collect 1 ml of blood in each tube. SHAKE tubes firmly. Do not centrifuge.					

Diagnostic Services Pty Ltd • ABN 26 009 503 100 • A subsidiary of Sonic Healthcare Limited

Redrafted May 2019

For all other tests please refer to our webpage with link to test collection manual, or contact the Laboratory

							,	
Approver	Approved .	16/05/2019	Owner	Michelle Leaman	Effective	16/05/2019	DSPL-SC-BF-0103-00.1	P.1 of 1